

Thermostable Enzymes for Clinical Chemistry

NIPRO Enzymes



From *Zymomonas mobilis*

ALCOHOL DEHYDROGENASE (ZM-ADH)
GLUCOKINASE (ZM-GlcK)
GLUCOSE-6-PHOSPHATE DEHYDROGENASE (ZM-G6PDH)

From *Bacillus stearothermophilus*

ACETATE KINASE (AK)
ADENYLATE KINASE (AdK)
ALANINE DEHYDROGENASE (AlaDH)
ALANINE RACEMASE (AlaR)
LEUCINE DEHYDROGENASE (LeuDH)
PHOSPHOGLUCOSE ISOMERASE (PGI)
PHOSPHOTRANSACETYLASE (PTA)
PYRUVATE KINASE (PK)
SUPEROXIDE DISMUTASE (SOD)

From Others

ACETATE KINASE (AK2)
BILIRUBIN OXIDASE (BOD3)
DIAPHORASE3 (DI-3)
DIAPHORASE22 (Di-22)
GALACTOSE DEHYDROGENASE (GalDH)
GLUCOKINASE2 (GlcK2)
GLUCOSE DEHYDROGENASE (GlcDH2)
D-LACTATE DEHYDROGENASE (D-LDH)
MALATE DEHYDROGENASE (MDH)
MUTAROTASE (MRO)
PHENYLALANINE DEHYDROGENASE (PheDH)
PHOSPHOTRANSACETYLASE (PTA2)
POLYNUCLEOTIDE PHOSPHORYLASE (PNPase3)
PYRUVATE KINASE (PK2)

ご照会は下記へお願い申し上げます。

ニプロ株式会社

本社/〒566-8510 大阪府摂津市千里丘新町3番26号

Tel 06-6373-3168

e-mail : nipro-web@nipro.co.jp

<http://www.nipro.co.jp/ja/>

For more information, please contact

NIPRO CORPORATION

3-26, Senriokashinmachi, Settsu,

Osaka, 566-8510, Japan

Phone +81 6 6373 3168

Fax +81 6 6373 8978

Bacillus stearothermophilus is used as a synonym of *Geobacillus stearothermophilus*.

2023.06

Quality

The Quality Management System of Enzyme Center, NIPRO Corp. has been certified as to meet the requirements of ISO9001 in the scope of design, development and manufacture of enzymes for analytical reagents and industrial use by JAPAN CHEMICAL QUALITY ASSURANCE LTD.

ALCOHOL DEHYDROGENASE (ZM-ADH)

[EC 1 .1 .1 .1]

from *Zymomonas mobilis*

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 400 U/mg protein	
Contaminants	: (as ZM-ADH activity = 100 %)	
	Glucose-6-phosphate dehydrogenase	< 0.10 %
	Glucokinase	< 0.02 %
	Pyruvate kinase	< 0.02 %
	NADH oxidase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 148,000	
Subunit molecular weight	: ca. 37,000	
Optimum pH	: 9.5 - 10.0	(Fig. 1)
pH stability	: 7.0 - 9.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 40 °C.	(Fig. 3, 4)
Michaelis constants	: (100 mM Glycine-KOH buffer, pH 9.0, at 30 °C)	
	Ethanol	110 mM
	Methanol	350 mM
	NAD ⁺	0.12 mM
	Acetaldehyde	1.66 mM
	NADH	0.03 mM
Substrate specificity	: Ethanol	100 %
	Methanol	0.05 %
	n-Propanol	42.3 %
	n-Butanol	0.28 %

STORAGE

Stable at -20 °C for at least six months


APPLICATION

The enzyme is useful for determination of alcohols or aldehydes.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of ZM-ADH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 80 mM Glycine-KOH, pH 9.5
- II NAD⁺ solution ; 10 mM (0.0663 g NAD⁺ free acid/10 mL distilled water)
- III Ethanol solution ; Ethanol (96 %)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM Tris succinate buffer containing 1mg/mL BSA and 0.2 mM CoCl₂, pH 7.0

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.
 - Solution I 22.90 mL
 - Solution II 6.00 mL
 - Solution III 1.10 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Neale, A.D., Scopes. R.K., Kelly, J.M., and Wettenhall, R.E.H.; *Eur. J. Biochem.*, **154**, 119 (1986)




Fig. 1 pH profile

(○ phosphate, ● Tris-HCl, ■ Gly-KOH)




Fig. 2 pH stability

treated for 24 hr at 4 °C in the following buffer solution (0.1 M), containing 0.5 mM CoCl₂;
 △ acetate, ○ phosphate, ● Tris-HCl, ■ Gly-KOH




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M phosphate buffer containing 0.2 mM CoCl₂, pH 6.5)





Fig. 4 Thermal stability

(treated in 0.1 M phosphate buffer containing 0.2 mM CoCl₂, pH 6.5
 ○ 50 °C, □ 55 °C, ● 60 °C)

GLUCOKINASE (ZM-GlcK)

[EC 2. 7. 1. 2]

from *Zymomonas mobilis*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 150 U/mg protein	
Contaminants	: (as ZM-GlcK activity = 100 %)	
	Glucose-6-phosphate dehydrogenase	< 0.02 %
	Phosphoglucomutase	< 0.01 %
	6-Phosphogluconate dehydrogenase	< 0.01 %
	Hexose-6-phosphate isomerase	< 0.01 %
	Glutathione reductase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 66,000	
Subunit molecular weight	: ca. 33,000	
Optimum pH	: 7.0 - 8.0	(Fig. 1)
pH stability	: 6.0 - 8.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 40 °C.	(Fig. 3, 4)
Michaelis constants	: (60 mM Phosphate buffer, pH 7.0, at 30 °C)	
	Glucose	0.10 mM
	ATP	0.65 mM
Activator	: Pi	

STORAGE

Stable at -20 °C for at least one year

APPLICATION


The enzyme is useful for diagnostic reagent, for example, glucose determination or CK determination, and for the specific determination of glucose.

Tris-HCl buffer is not suitable for the practical use of ZM-GlcK.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of ZM-GlcK that forms 1 μmol of glucose-6-phosphate per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Triethanolamine - NaOH and 3 mM K_2HPO_4 , pH 7.5
- II ATP solution ; 100 mM (0.605 g ATP disodium salt·3H₂O/(8.2 mL distilled water + 1.8 mL 1 N-NaOH))
- III MgCl_2 solution ; 1 M (20.33 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ /100 mL distilled water)
- IV NAD^+ solution ; 100 mM (0.663 g NAD^+ free acid/10 mL distilled water)
- V Glucose solution ; 40mM (0.072 g glucose (anhyd.)/10 mL distilled water)
- VI Glucose-6-phosphate dehydrogenase (G6PDH) ; 2000 U/mL (from *Zymomonas mobilis*, Nipro Corp., Dissolve with Buffer solution I)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM potassium phosphate buffer containing 1 mg/mL BSA, pH 7.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	20.07 mL	Solution IV	0.60 mL
Solution II	1.50 mL	Solution V	7.50 mL
Solution III	0.30 mL	Solution VI	0.03 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Scopes. R.K., Testolin, V., Stoter, A., Griffiths-Smith, K., and Algar, E.M.; *Biochem. J.*, **228**, 627 (1985)




Fig. 1 pH profile

(○ MES-KOH, □ TEA-NaOH,
■ Gly-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
following buffer solution (0.1 M);
△ acetate, ○ phosphate,
● Tris-HCl, ■ Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
phosphate buffer, pH 7.0)




Fig. 4 Thermal stability

(treated in 0.1 M phosphate
buffer, pH 7.0
○ 40 °C, □ 50 °C, ● 60 °C)

GLUCOSE-6-PHOSPHATE DEHYDROGENASE (ZM-G6PDH)

[EC 1. 1. 1. 49]

from *Zymomonas mobilis*

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 250 U/mg protein	
Contaminants	: (as ZM-G6PDH activity = 100 %)	
	Glucokinase	< 0.02 %
	Phosphoglucomutase	< 0.01 %
	6-Phosphogluconate dehydrogenase	< 0.02 %
	Hexose-6-phosphate isomerase	< 0.01 %
	Glutathione reductase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 208,000	
Subunit molecular weight	: ca. 52,000	
Optimum pH	: 8.0	(Fig. 1)
pH stability	: 5.0 - 10.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 50 °C.	(Fig. 3, 4)
Michaelis constants	: (30 mM Tris-HCl buffer, pH 8.0, at 30 °C)	
	Glucose-6-phosphate	0.14 mM
	NADP ⁺	0.02 mM
	NAD ⁺	0.14 mM
Substrate specificity	: NADP ⁺	70 %
	NAD ⁺	100 %

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for diagnostic reagent, for example, glucose determination or CK determination, and for the specific determination of glucose.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of ZM-G6PDH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 50 mM Tris-HCl, pH 8.0
- II NAD⁺ solution ; 100 mM (0.663 g NAD⁺ free acid/10 mL distilled water)
- III Glucose-6-phosphate (G6P) solution ; 33 mM (0.112 g G6P disodium salt 2H₂O/10 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM potassium phosphate buffer containing 1 mg/mL BSA, pH 7.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.
 - Solution I 26.40 mL
 - Solution II 0.90 mL
 - Solution III 2.70 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Scopes, R.K., Testolin, V., Stoter, A., Griffiths-Smith, K., and Algar. E.M.; *Biochem. J.*, **228**. 627 (1985)




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M);
 Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
 phosphate buffer, pH 7.0)




Fig. 4 Thermal stability

(treated in 0.1 M phosphate
 buffer, pH 7.0
 \circ 50 °C, \square 55 °C, \bullet 60 °C)

ACETATE KINASE (AK)

[EC 2. 7. 2. 1]

from *Bacillus stearothermophilus*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 1,100 U/mg protein	
Contaminants	: (as AK activity = 100 %)	
	Lactate dehydrogenase	< 0.01 %
	Adenylate kinase	< 0.01 %
	NADH oxidase	< 0.01 %
	GOT	< 0.01 %
	GPT	< 0.01 %

PROPERTIES

Molecular weight	: ca. 160,000	
Subunit molecular weight	: ca. 40,000	
Optimum pH	: 7.2	(Fig. 1)
pH stability	: 7.0 - 8.0	(Fig. 2)
Isoelectric point	: 4.8	
Thermal stability	: No detectable decrease in activity up to 65 °C.	(Fig. 3, 4)
Michaelis constants	: (57 mM Imidazole- HCl buffer, pH 7.2, at 30 °C)	
	Acetate	120 mM
	Acetylphosphate	2.3 mM
	ATP	1.2 mM
	ADP	0.8 mM
Substrate specificity	: Acetate	100 %
	Formate	0 %
	Propionate	5 %
	Butyrate	0 %
	Oxalate	0 %
	Citrate	0 %
	Malate	0 %
	Glycine	0 %
Activator	: Fructose-1,6-bisphosphate	

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for determination of acetate or for ATP regeneration system.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of AK that forms 1 μmol of ADP per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Imidazole-HCl, pH 7.2
- II ATP solution ; 100 mM (0.605 g ATP disodium salt·3H₂O/(8.2 mL distilled water + 1.8 mL 1 N-NaOH))
- III Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- IV NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)
- V MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O /100 mL distilled water)
- VI KCl solution ; 2.5 M (18.64 g KCl/100 mL distilled water)
- VII Pyruvate kinase (PK) ; (from rabbit muscle, Roche Diagnostics K.K., No. 128 155) crystalline suspension in 3.2 M (NH₄)₂SO₄ solution (10 mg/mL) approx. 200 U/mg at 25 °C
- VIII Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., Product Code: LDH-02) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C
- IX Sodium acetate solution ; 2 M (27.22 g sodium acetate·3H₂O/100 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM potassium phosphate buffer, pH 7.5.

Procedure

1. Prepare the following reaction mixture and pipette 2.4 mL of reaction mixture into a cuvette.

Solution I	16.92 mL	Solution V	0.60 mL
Solution II	3.00 mL	Solution VI	0.90 mL
Solution III	1.80 mL	Solution VII	0.12 mL
Solution IV	0.60 mL	Solution VIII	0.06 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.60 mL of Solution IX and 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH (cm²/μmol)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Nakajima, H., Suzuki, K., and Imahori, K. ; *J. Biochem.*, **84**, 193 (1978)
2. Nakajima, H., Suzuki, K., and Imahori, K. ; *ibid.*, **84**, 1139 (1978)
3. Nakajima, H., Suzuki, K., and Imahori, K. ; *ibid.*, **86**, 1169 (1979)




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
● Tris-HCl)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
following buffer solution (0.1 M);
 \circ phosphate, ● Tris-HCl,
▲ carbonate)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
potassium phosphate buffer,
pH 7.5)




Fig. 4 Thermal stability

(treated in 0.1M potassium
phosphate buffer, pH 7.5
 \circ 60 °C, \square 65 °C, ● 70 °C)

ADENYLATE KINASE (AdK)

[EC 2. 7. 4. 3]

from *Bacillus stearothermophilus*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 200 U/mg protein	
Contaminants	: (as AdK activity = 100 %)	
	ATPase	< 0.01 %
	Phosphoglycerate kinase	< 0.10 %

PROPERTIES

Molecular weight	: ca. 20,000	
Optimum pH	: 6.5	(Fig. 1)
pH stability	: 8.0 - 10.5	(Fig. 2)
Isoelectric point	: 5.0	
Thermal stability	: No detectable decrease in activity up to 65 °C.	(Fig. 3, 4)
Michaelis constants	: (89 mM Imidazole-HCl buffer, pH 6.5, at 30 °C)	
	ATP	0.04 mM
	ADP	0.05 mM
	AMP	0.02 mM

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for determination of AMP or for system involving ATP regeneration.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of AdK that forms 2 μmol of ADP per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Imidazole-HCl, pH 6.5
- II AMP solution ; 50 mM (0.250 g AMP disodium salt·6H₂O/10 mL distilled water)
- III ATP solution ; 100 mM (0.605 g ATP disodium salt·3H₂O/(8.2 mL distilled water + 1.8 mL 1 N-NaOH))
- IV NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O /10 mL distilled water)
- V Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- VI MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O/100 mL distilled water)
- VII KCl solution ; 2.5 M (18.64 g KCl/100mL distilled water)
- VIII Pyruvate kinase (PK) ; (from rabbit muscle, Roche Diagnostics K.K., No. 128 155) crystalline suspension in 3.2 M (NH₄)₂SO₄ solution (10 mg/mL) approx. 200 U/mg at 25 °C
- IX Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., Product Code: LDH-02) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 2.5 to 5 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	26.70 mL	Solution VI	0.60 mL
Solution II	0.24 mL	Solution VII	1.20 mL
Solution III	0.30 mL	Solution VIII	0.09 mL
Solution IV	0.60 mL	Solution IX	0.09 mL
Solution V	0.18 mL		
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{2 \times 6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

2 ; according to the reaction that forms 2 μmol of ADP, one unit of activity of Adk is defined to form 2 μmol of ADP.

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Imahori, K., Nakajima, H., Nagata, K., and Iwasaki, T.; *Seikagaku*, **53**, 829 (1981)




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
● Tris-HCl, ▲ carbonate)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
following buffer solution (0.1 M);
 Δ acetate, \circ phosphate,
● Tris-HCl, ▲ carbonate)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
Tris-HCl buffer, pH 9.0)




Fig. 4 Thermal stability

(treated in 0.1 M Tris-HCl buffer,
pH 9.0
 \circ 60 °C, \square 65 °C, \bullet 70 °C)

ALANINE DEHYDROGENASE (AlaDH)

[EC 1. 4. 1. 1]

from *Bacillus stearothermophilus*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 10 U/mg protein	
Contaminants	: (as AlaDH activity = 100 %)	
	NADH oxidase	< 0.01 %
	Lactate dehydrogenase	< 0.10 %

PROPERTIES

Molecular weight	: ca. 230,000	
Subunit molecular weight	: ca. 38,000	
Optimum pH	: 10.4	(Fig. 1)
pH stability	: 7.0 - 11.5	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 70 °C.	(Fig. 3, 4)
Michaelis constants	: (125 mM Glycine-NaOH buffer, pH 10.5, at 30 °C)	
	L-Alanine	10.0 mM
	NAD ⁺	0.26 mM
Substrate specificity	: L-Alanine	100 %
	L-Leucine	0 %
	L-Isoleucine	0 %

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for determination of L-alanine.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of AlaDH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 250 mM Glycine-NaOH, pH 10.5
- II L-Alanine solution ; 150 mM (1.336 g L-alanine/80 mL distilled water, adjusted to pH 10.5 with 1 N-NaOH and filled up to 100 mL with distilled water)
- III NAD^+ solution ; 100 mM (0.663 g NAD^+ / 10 mL with distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 100 mM glycine - NaOH buffer, pH 9.5.

Procedure

- Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	15.00 mL	Solution III	1.50 mL
Solution II	10.00 mL	H ₂ O	3.50 mL
- Incubate at 30 °C for about 3 minutes.
- Add 0.01 mL of enzyme solution into the cuvette and mix.
- Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

- Sakamoto, Y., Nagata, S., Esaki, N., Tanaka, H. and Soda, K.; *J. Ferment. Bioeng.*, **69**, 154 (1990)




Fig. 1 pH profile

(● Tris-HCl, ■ Gly-KOH, ○ phosphate)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
△ acetate, ○ phosphate,
● Tris-HCl, ■ Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M Gly-KOH buffer, pH 9.0)




Fig. 4 Thermal stability

(treated in 0.1 M Gly-KOH buffer, pH 9.0
○ 70 °C, □ 80 °C, ● 90 °C)

ALANINE RACEMASE (AlaR)

[EC 5. 1. 1. 1]

from *Bacillus stearothermophilus*

D-Alanine ↔ L-Alanine

SPECIFICATION

State	: Liquid	
Specific activity	: more than 950 U/mg protein	
Contaminants	: (as AlaR activity = 100 %)	
	Lactate dehydrogenase	< 0.01 %
	NADH oxidase	< 0.01 %
	Alanine dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 78,000	
Subunit molecular weight	: ca. 39,000	
Optimum pH	: 10.5 - 12.0	(Fig. 1)
pH stability	: 5.5 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 70 °C.	(Fig. 3, 4)
Michaelis constants	: (100 mM Carbonate buffer, pH 10.5, at 30 °C)	
	D-Alanine	31 mM
Substrate specificity	:	


STORAGE

Stable at least one year at -25 °C.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of AlaR that forms 1 μmol of L-alanine per minute at 30 °C.

Solutions

- I Buffer solution ; 200 mM Sodium hydrogencarbonate, pH 10.5
- II D-Alanine solution ; 1 M (0.891 g D-alanine/10 mL distilled water)
- III NAD^+ solution ; 100 mM (0.663 g NAD^+ /10 mL distilled water)
- IV L-Alanine dehydrogenase (AlaDH) ; 1000 U/mL (from *Bacillus stearothermophilus*, Nipro Corp., Dissolve with distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM potassium phosphate buffer, pH 7.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	16.50 mL	Solution IV	1.50 mL
Solution II	3.00 mL	H_2O	8.25 mL
Solution III	0.75 mL		
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Inagaki, K., Tanizawa, K., Badet, B., Walsh, C.T., Tanaka, H., and Soda, K.; *Biochemistry*, **25**, 3268 (1986)




Fig. 1 pH profile

(○ phosphate, ● Tris-HCl, ■ Gly-KOH, □ NaHCO₃-NaOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following buffer solution (0.2 M);
△ acetate, ○ phosphate, ● Tris-HCl, ■ Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 50 mM Tris-HCl buffer, pH 9.0)




Fig. 4 Thermal stability

(treated in 50 mM Tris-HCl buffer, pH 9.0
○ 70 °C, □ 80 °C, ● 85 °C)




Fig. 5 Stability (Liquid form) at -25 °C

LEUCINE DEHYDROGENASE (LeuDH)

[EC 1. 4. 1. 9]

from *Bacillus stearothermophilus*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 40 U/mg protein	
Contaminants	: (as LeuDH activity = 100 %)	
	NADH oxidase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 300,000	
Subunit molecular weight	: ca. 49,000	
Optimum pH	: 10.6	(Fig. 1)
pH stability	: 6.0 - 11.5	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)
Michaelis constants	: (125mM Sodium phosphate buffer, pH 10.5, at 30 °C)	
	L-Leucine	3.4 mM
	NAD ⁺	0.3 mM
Substrate specificity	: L-Leucine	100 %
	L-Valine	86 %
	L-Isoleucine	73 %

STORAGE

Stable at -20 °C for at least one year

APPLICATION

The enzyme is useful for determination of L-leucine, L-valine or L-isoleucine.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of LeuDH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 250 mM Sodium phosphate, pH 10.5
- II L-Leucine solution ; 60 mM (0.787 g L-leucine/80 mL distilled water, adjusted to pH 10.5 with 1 N-NaOH and filled up to 100 mL with distilled water)
- III NAD^+ solution ; 100mM (0.663 g NAD^+ / 10mL with distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 100 mM sodium phosphate buffer, pH 9.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette,

Solution I	15.00 mL	Solution III	0.93 mL
Solution II	10.00 mL	H_2O	4.07 mL
2. Incubate at 30°C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Ohshima, T., Nagata, S., and Soda, K.; *Arch. Microbiol.*, **141**, 407 (1985)




Fig. 1 pH profile

(■ Gly-KOH, ○ phosphate)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following
buffer solution (0.1 M);
△ acetate, ○ phosphate,
● Tris-HCl, ■ Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M Gly-KOH buffer, pH 9.0)




Fig. 4 Thermal stability

(treated in 0.1M Gly-KOH buffer, pH 9.0
○ 60 °C, □ 70 °C, ● 80 °C)

PHOSPHOGLUCOSE ISOMERASE (PGI)

[EC 5. 3. 1. 9]

from *Bacillus stearothermophilus*

D-Glucose-6-phosphate ↔ D-Fructose-6-phosphate

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 400 U/mg protein	
Contaminants	: (as PGI activity = 100 %)	
	Phosphofructokinase	< 0.01 %
	6-Phosphogluconate dehydrogenase	< 0.01 %
	Phosphoglucomutase	< 0.01 %
	NADPH oxidase	< 0.01 %
	Glutathione reductase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 200,000	
Subunit molecular weight	: ca. 54,000	
Optimum pH	: 9.0 - 10.0	(Fig. 1)
pH stability	: 6.0 - 10.5	(Fig. 2)
Isoelectric point	: 4.2	
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)
Michaelis constants	: (95 mM Tris-HCl buffer, pH 9.0, at 30 °C)	
	Fructose-6-phosphate	0.27 mM

STORAGE

Stable at -20 °C for at least one year

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of PGI that forms 1 μmol of glucose-6-phosphate per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH 9.0
- II Fructose-6-phosphate (F6P) solution ; 100 mM (0.310 g F6P disodium salt/10 mL distilled water)
- III NADP⁺ solution ; 22.5 mM (0.188 g NADP⁺ sodium salt·4H₂O/10 mL distilled water)
- IV Glucose-6-phosphate dehydrogenase (G6PDH) ; (from yeast, Roche Diagnostics K.K., No. 127 671) suspension in 3.2 M (NH₄)₂SO₄ solution (10 mg/2 mL) approx. 140 U/mg at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	28.44 mL	Solution III	0.60 mL
Solution II	0.90 mL	Solution IV	0.06 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of the curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADPH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Muramatsu, N., and Nosoh, T.; *Arch. Biochem. Biophys.*, **144**, 245 (1971)




Fig. 1 pH profile

(
 △ acetate, ○ phosphate,
 ● Tris-HCl, ■ Gly-KOH
)




Fig. 2 pH stability

(
 treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M);
 △ acetate, ○ phosphate,
 ● Tris-HCl, ■ Gly-KOH
)




Fig. 3 Thermal stability

(
 treated for 15 min in 50 mM
 Tris-HCl buffer, pH 8.5
)




Fig. 4 Thermal stability

(
 treated in 50 mM Tris-HCl
 buffer, pH 8.5
 ○ 60 °C, □ 65 °C, ● 70 °C
)

PHOSPHOTRANSACETYLASE (PTA)

[EC 2. 3. 1. 8]

from *Bacillus stearothermophilus*

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 5,000 U/mg protein	
Contaminants	: (as PTA activity = 100 %)	
	Acetate kinase	< 0.01 %
	Adenylate kinase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 70,000	
Subunit molecular weight	: ca. 35,000	
Optimum pH	: 7.5	(Fig. 1)
pH stability	: 7.0 - 11.0	(Fig. 2)
Isoelectric point	: 4.5	
Thermal stability	: No detectable decrease in activity up to 50 °C.	(Fig. 3, 4)
Michaelis constants	: (87 mM Tris-HCl buffer, pH 7.5, at 30 °C)	
	Coenzyme A	0.4 mM
	Acetyl Phosphate	1.1 mM

STORAGE

Stable at -20 °C for at least one year

APPLICATION

The enzyme is useful for determination of CoA or acetate.

ASSAY

Principle

The change in absorbance is measured at 233 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of PTA that forms 1 μmol of acetyl-CoA per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH 7.5
- II CoA solution ; 6.4 mM (50 mg CoA trilithium salt/10 mL distilled water)
- III Acetylphosphate solution ; 217 mM (0.400 g acetylphosphate potassium lithium salt/10 mL distilled water)
- IV Ammonium sulfate (AmS) solution ; 1 M (13.2 g AmS/100 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 20 U/mL with 50 mM Tris-HCl buffer, pH 8.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	26.0 mL	Solution III	1.0 mL
Solution II	2.0 mL	Solution IV	1.0 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 233 nm per minute (ΔAbs_{233}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{233}) \times (3.00 + 0.01)}{4.44 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

4.44 ; differential millimolar extinction coefficient between acetyl-CoA and CoA ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method




Fig. 1 pH profile
 [● Tris-HCl]




Fig. 2 pH stability
 (treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
 △ acetate, ○ phosphate,
 ● Tris-HCl, ■ Gly-KOH)




Fig. 3 Thermal stability
 (treated for 15 min in 50 mM Tris-HCl buffer, pH 8.0)




Fig. 4 Thermal stability
 (treated in 50 mM Tris-HCl buffer, pH 8.0
 ○ 50 °C, □ 60 °C, ● 65 °C)

PYRUVATE KINASE (PK)

[EC 2.7.1.40]

from *Bacillus stearothermophilus*

ATP + Pyruvate \leftrightarrow ADP + Phosphoenolpyruvate

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 230 U/mg protein	
Contaminants	: (as PK activity = 100 %)	
	Adenylate kinase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 260,000	
Subunit molecular weight	: ca. 68,000	
Optimum pH	: 7.0	(Fig. 1)
pH stability	: 8.0 - 10.0	(Fig. 2)
Isoelectric point	: 5.2	
Thermal stability	: No detectable decrease in activity up to 55 °C.	(Fig. 3, 4)
Michaelis constants	: (76 mM Imidazole-HCl buffer, pH 7.2, at 30 °C)	
	Phosphoenolpyruvate	0.6 mM
	ADP	0.9 mM

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for diagnostic reagent, for example, ADP determination.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of PK that forms 1 μmol of pyruvate per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Imidazole-HCl, pH 7.2
- II ADP solution ; 100 mM (0.507 g ADP disodium salt·2H₂O/(9.0 mL distilled water + 1.0 mL 1 N NaOH))
- III NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)
- IV Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- V MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O/100 mL distilled water)
- VI KCl solution ; 2.5 M (18.64 g KCl/100 mL distilled water)
- VII Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., Product Code: LDH-02) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	22.71 mL	Solution V	0.48 mL
Solution II	2.40 mL	Solution VI	0.90 mL
Solution III	0.45 mL	Solution VII	0.06 mL
Solution IV	3.00 mL		
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH (cm²/μmol)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Sakai, H., Suzuki, K., and Imahori, K.; *J. Biochem.*, **99**, 1157 (1986)




Fig. 1 pH profile

(Δ acetate, \square imidazole-HCl, \bullet Tris-HCl)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
 Δ acetate, \square imidazole-HCl, \bullet Tris-HCl, \blacktriangle carbonate)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M Tris-HCl buffer, pH 8.5)





Fig. 4 Thermal stability

(treated in 0.1 M Tris-HCl buffer, pH 8.5
 \circ 55 °C, \square 60 °C, \bullet 65 °C)

SUPEROXIDE DISMUTASE (SOD)

[EC 1.15.1.1]

from *Bacillus stearothermophilus*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 9,000 U/mg protein	
Contaminants	: (as SOD activity = 100 %)	
	Catalase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 50,000	
Subunit molecular weight	: ca. 25,000	
Metal content	: 1.5 g atoms of Mn per mole of enzyme	
Optimum pH	: 9.5	(Fig. 1)
pH stability	: 6.0 - 9.0	(Fig. 2)
Isoelectric point	: 4.5	
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for medicine, cosmetic material and nutrition or antioxidant.

ASSAY

Principle

To determine the enzyme activity of cytochrome c reduction is measured by the following reactions.



Unit Definition

One unit of activity is defined as the amount of SOD required to inhibit the rate of reduction of cytochrome C by 50 % at 30 °C.

Solutions

- I Buffer solution ; 75 mM Potassium phosphate buffer, pH 7.8
- II Xanthine solution ; 0.75 mM (0.010 g xanthine/50 mL N/250 NaOH)
- III Cytochrome c solution ; 0.15 mM (0.019 g cytochrome c/10 mL distilled water, Sigma-Aldrich Co., No. C-2506, from horse heart)
- IV EDTA solution ; 1.5 mM (0.028 g EDTA disodium salt·2H₂O/50 mL distilled water)
- V Xanthine oxidase (XOD) ; (from buttermilk, Sigma-Aldrich Co., No. X-1875) suspension in 2.3 M (NH₄)₂SO₄ solution is diluted to 0.04 U/mL with distilled water. (prepare freshly)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to approx. 600 U/mL with 50 mM potassium phosphate buffer, pH 7.5.

Procedure

1. Prepare the following reaction mixture and pipette 2.80 mL of reaction mixture and 0.005 mL of enzyme solution into a cuvette.

Solution I	22.00 mL	Solution III	2.00 mL
Solution II	2.00 mL	Solution IV	2.00 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.20 mL of Solution V into the cuvette and mix.
4. Read absorbance change at 550 nm per minute for the linear portion of curve (ΔAbs•test)*.
5. Add 0.005 mL of Solution I in place of enzyme solution and measure the same above 4 (ΔAbs•blank).

*Dilute enzyme solution with 50 mM potassium phosphate buffer, pH 7.5, because the decrease in the initial rate should not fall outside the range of 40 to 60 % for the results to be valid.

Calculation

$$\text{Volume activity (U/mL)} = \left[\frac{(\Delta\text{Abs}\cdot\text{blank})}{(\Delta\text{Abs}\cdot\text{test})} - 1 \right] \times \frac{601}{1} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{protein concentration (mg/mL)*}}$$

d.f. ; dilution factor

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Bridgen, J., Harris, J.I., and Kolb, E.; *J. Mol. Biol.*, **105**, 333 (1976)
2. Brock, C.J., Harris, J.I., and Sato, S.; *ibid.*, **107**, 175 (1976)
3. Brock, C.J., and Walker, J.E.; *Biochemistry*, **19**, 2873 (1980)
4. Auffret, A.D., Blake, T.J., and Williams, D.H.; *Eur. J. Biochem.*, **113**, 333 (1981)
5. Atkinson, T., Banks, G.T., Bruton, C.J., Comer, M.J., Jakes, R., Kamalagharan, T., Whitak, A.R., and Winter, G.P.; *J. Appl. Biochem.*, **1**, 247 (1979)




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
● Tris-HCl, \blacktriangle carbonate)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
Britton-Robinson buffer)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M potassium
phosphate buffer, pH 7.5)




Fig. 4 Thermal stability

(treated in 0.1 M potassium
phosphate buffer, pH 7.5
 \circ 60 °C, \square 70 °C, \bullet 80 °C)

ACETATE KINASE (AK2)

[EC 2. 7. 2. 1]

from recombinant *E. coli*

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 1,100 U/mg protein	
Contaminants	: (as AK2 activity = 100 %)	
	Lactate dehydrogenase	< 0.01 %
	Adenylate kinase	< 0.01 %
	NADH oxidase	< 0.01 %
	GOT	< 0.01 %
	GPT	< 0.01 %

PROPERTIES

Molecular weight	: ca. 172,000	
Subunit molecular weight	: ca. 39,400	
Optimum pH	: 6.0	(Fig. 1)
pH stability	: 6.0 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 65 °C.	(Fig. 3, 4)
Michaelis constants	: (57 mM Imidazole-HCl buffer, pH 7.2, at 30 °C)	
	Acetate	130 mM
	Acetylphosphate	3.3 mM
	ATP	5.4 mM
	ADP	0.6 mM
Substrate specificity	: Acetate	100 %
	Formate	0 %
	Propionate	4 %
	Butyrate	1 %
	Oxalate	0 %
	Malate	0 %
	Glycine	0 %
Activator	: Fructose-1,6-bisphosphate	


STORAGE

Stable at -20 °C for at least one year

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of AK that forms 1 μmol of ADP per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Imidazole-HCl, pH 7.2
- II ATP solution ; 100 mM (0.605 g ATP disodium salt·3H₂O/(8.2 mL distilled water + 1.8 mL 1 N-NaOH))
- III Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- IV NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)
- V MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O /100 mL distilled water)
- VI KCl solution ; 2.5 M (18.64 g KCl/100 mL distilled water)
- VII Pyruvate kinase (PK) ; (from rabbit muscle, Roche Diagnostics K.K., No. 128 155) crystalline suspension in 3.2 M (NH₄)₂SO₄ solution (10 mg/mL) approx. 200 U/mg at 25 °C
- VIII Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., Product Code: LDH-02) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C
- IX Sodium acetate solution ; 2 M (27.22 g sodium acetate·3H₂O/100 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM potassium phosphate buffer, pH 7.5.

Procedure

1. Prepare the following reaction mixture and pipette 2.4 mL of reaction mixture into a cuvette.

Solution I	16.92 mL	Solution V	0.60 mL
Solution II	3.00 mL	Solution VI	0.90 mL
Solution III	1.80 mL	Solution VII	0.12 mL
Solution IV	0.60 mL	Solution VIII	0.06 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.60 mL of Solution IX and 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH (cm²/μmol)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Nakajima, H., Suzuki, K., and Imahori, K. ; *J. Biochem.*, **84**, 193 (1978)
2. Nakajima, H., Suzuki, K., and Imahori, K. ; *ibid.*, **84**, 1139 (1978)
3. Nakajima, H., Suzuki, K., and Imahori, K. ; *ibid.*, **86**, 1169 (1979)




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M);
 Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacktriangle Carbonate)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
 phosphate buffer, pH 7.5)





Fig. 4 Thermal stability

(treated in 0.1 M phosphate
 buffer, pH 7.5
 \circ 60 °C, \square 65 °C, \bullet 70 °C)

BILIRUBIN OXIDASE (BOD3)

[EC 1.3.3.5]

from *Trachyderma tsunodae*



SPECIFICATION

State : Lyophilized
Specific activity : more than 100 U/mg protein

PROPERTIES

Molecular weight	: ca. 60,000 (SDS-electrophoresis)	
	: ca. 80,000 (Gel filtration)	
Optimum pH	: 5.0	(Fig. 1)
pH stability	: 4.0 – 11.0 (4 °C, 24 hr)	(Fig. 2)
Isoelectric point (calculation)	: 3.8	
Optimum temperature	: 65 – 80 °C	(Fig. 3)
Thermal stability	: No detectable decrease in activity up to 50 °C. (pH 7.0)	(Fig. 4, 5)
Michaelis constants	: See table 1	
Substrate specificity	: See table 1	

STORAGE

Stable at -20 °C for one year


APPLICATION

The enzyme is useful for enzymatic determination of bilirubin.
It could be used as a cathode catalyst in biofuel cells.

ASSAY

Principle

The change in absorbance is measured at 500 nm according to the following reaction.



Unit Definition

One unit of activity is defined according to the calculation formula below.

Solutions

- I Buffer solution ; 300 mM Potassium phosphate buffer, pH7.0
- II 4-Aminoantipyrine (4-AA) solution ; 24.6 mM (0.25 g 4-AA / 50 mL distilled water)
- III Phenol solution ; 420 mM (1.98 g phenol/50mL distilled water)
- IV Peroxidase*¹ (POD) solution ; 240 U/mL (2,400 U/10mL distilled water)

*¹POD: TOYOBO Co., LTD. Grade III #PEO-302

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 15 to 60 U/mL with 10 mM potassium phosphate buffer, pH 7.0 containing 0.1 % BSA.

Procedure

1. Prepare the following reaction mixture and pipette 0.90 mL of reaction mixture into a cuvette.

Solution I	4.00 mL
Solution II	0.40 mL
Solution III	0.40 mL
Solution IV	0.40 mL
H ₂ O	6.40 mL
2. Incubate at 37 °C for about 3 minutes.
3. Add 0.005 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 500 nm per minute (ΔAbs (test)) in linear portion of curve. Repeat the procedure 3 using distilled water in place of enzyme solution, and ΔAbs (blank) is obtained.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs (test)} - \Delta\text{Abs (blank)}) \times (0.90 + 0.005)}{11.11 \times 0.005 \times 1/20} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^{*2}}$$

d.f. ; dilution factor

11.11 ; millimolar extinction coefficient of quinoneimine dye at 500 nm (cm²/μmol)

1/20 ; coefficient of transformation for internal unit definition

*²Protein concentration ; determined by Bradford's method




Fig. 1 pH profile

△ acetate, ○ phosphate
 ● Tris-HCl, ■ Glycine-KOH




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (50 mM);
 △ acetate, ○ phosphate
 ● Tris-HCl, ■ Glycine-KOH)




Fig. 3 Thermal activity




Fig. 4 Thermal stability

(treated for 15 min in 20 mM
 potassium phosphate buffer, pH 7.0)




Fig. 5 Thermal stability

treated in 20 mM potassium phosphate buffer, pH 7.0
 ○ 50 °C, ● 55 °C, □ 60 °C, ■ 65 °C




Fig. 6 pH profile (ABTS*³)




Fig. 7 pH profile (Bilirubin C*⁴)




Fig. 8 pH profile (Bilirubin F*⁴)

Measured in 20 mM buffer.
 ▲ Glycine-HCl, △ acetate, ○ phosphate, ● Tris-HCl, ■ Glycine-KOH

*³ 2,2'-Azinobis (3-ethylbenzothiazoline-6-sulfonic Acid) Diammonium Salt

*⁴ Bilirubin C (conjugated type) and Bilirubin F (free type) are from [Interference Check. A Plus] (Sysmex, Kobe, Japan).

Table 1. Substrate specificity of BOD3


	Phenol	ABTS	Bilirubin C	Bilirubin F
Optimum pH	5.0	4.0	6.0	6.0
Michaelis constants (μM)	41	39	26	26
Relative activity (%)	100	427	36	8
Wavelength for Measurement (nm)	500	405	450	450
Extinction Coefficient ($\text{cm}^2/\mu\text{mol}$)	11.11	29	74	32

Michaelis constants and activity of phenol were defined at pH 7.0. They were defined at each optimum pH when the substrate was ABTS, Bilirubin C, or Bilirubin F,.

DIAPHORASE 3 (DI-3)

[EC 1. 6. 99. -]

from recombinant *E. coli*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 1,000 U/mg protein	
Contaminants	: (as Diaphorase activity = 100 %)	
	Adenylate kinase	< 0.01 %
	NADH oxidase	< 0.01 %

PROPERTIES

Subunit molecular weight	: ca. 20,000	(SDS-electrophoresis)	
Optimum pH	: 8.0		(Fig. 1)
pH stability	: 7.5 - 9.5		(Fig. 2)
Isoelectric point	: 4.7		
Thermal stability	: No detectable decrease in activity up to 60 °C.		(Fig. 3, 4)
Michaelis constants	: See Table 1		

STORAGE

Stable at -20 to 5 °C for one year


APPLICATION

The enzyme is useful for the measurement of various dehydrogenase reactions in visible spectral range.

ASSAY

Principle

The change in absorbance is measured at 600 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of DI-3 that reduces 1 μmol of DCIP per minute at 30 °C.

Solutions

- I Buffer solution ; 500 mM Tris-HCl, pH 8.5
- II NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)
- III 2,6-Dichlorophenolindophenol (DCIP) solution ; 1.2 mM (2.0 mg DCIP sodium salt·2H₂O/5mL distilled water) (prepare freshly)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 1.0 to 2.0 U/mL with 50 mM potassium phosphate buffer, pH 7.5.

Procedure

1. Prepare the following reaction mixture and pipette 2.28 mL of reaction mixture and 0.12 mL of Solution III into a cuvette.

Solution I	3.00 mL
Solution II	2.28 mL
H ₂ O	23.22 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.008 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 600 nm per minute ($\Delta\text{Abs}(\text{test})$) in linear portion of curve. Repeat the Procedure 3 using distilled water in place of enzyme solution, and $\Delta\text{Abs}(\text{blank})$ is obtained.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs (test)} - \Delta\text{Abs (blank)}) \times (2.40 + 0.008)}{19 \times 0.008} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

19 ; millimolar extinction coefficient of DCIP ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Mains, I., Power, D.M., Thomas, E.W. and Buswell J. A.; *Biochem. J.*, **191**, 457 (1980)




Fig. 1 pH profile

○ phosphate, ● Tris-HCl,
■ Gly-KCl-KOH




Fig. 2 pH stability

treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
△ acetate, ○ phosphate,
● Tris-HCl




Fig. 3 Thermal stability

treated for 15 min in 0.1 M potassium phosphate buffer, pH 7.5




Fig. 4 Thermal stability

treated in 0.1 M potassium phosphate buffer, pH 7.5
○ 50 °C, □ 60 °C, ● 70 °C

Table 1. SUBSTRATE SPECIFICITY OF DIAPHORASE

Acceptor	DCIP* ¹	NTB* ²	MTT* ³
K _m ^{Acceptor} (mM)	0.02	0.15	0.9
K _m ^{NADH} (mM)	0.37	0.01	0.05
K _m ^{NADPH} (mM)	32.7	0.31	2.0
Optimum pH	8.0	10	8.0
Assay Mixture	Tris-HCl (pH 8.5) 50 mM NAD(P)H 1 mM DCIP 0.06 mM	TEA (pH 7.0) 50 mM NAD(P)H 1 mM NBT 0.5 mM Triton X-100 0.1 %	TEA (pH 7.0) 50 mM NAD(P)H 1 mM MTT 0.5 mM Triton X-100 0.5 %
Wavelength for Measurement (nm)	600	550	565
Extinction Coefficient (cm ² /μmol)	19	12.4	20

*1 2,6-Dichlorophenolindophenol

*2 Nitrotertrazolium Blue

*3 Thiazolyl Blue Tetrazolium Bromide

pH profiles of DI-3 (Acceptor; NTB or MTT)




Fig. 5 pH profile (NTB)

□ triethanolamine, ○ phosphate,
● Tris-HCl, ■ Gly-KCl-KOH





Fig. 6 pH profile (MTT)

□ triethanolamine, ○ phosphate,
● Tris-HCl, ■ Gly-KCl-KOH

DIAPHORASE 22 (Di-22)

[EC 1. 8. 1. 4]

from recombinant *E.coli*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 150 U/mg protein	
Contaminants	: (as Diaphorase activity = 100 %)	
	Adenylate kinase	< 0.01 %
	NADH oxidase	< 0.20 %

PROPERTIES

Molecular weight	: ca. 110,000	
Subunit molecular weight	: ca. 50,000	
Optimum pH	: 8.0	(Fig,1)
pH stability	: 6.0 – 9.0	(Fig.2)
Thermal stability	: No detectable decrease in activity up to 70 °C.	(Fig. 3, 4)
Michaelis constants	: (50 mM HEPES buffer, pH 7.0, at 30 °C)	
	3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2 <i>H</i> -tetrazolium bromide (MTT)	0.345 mM
	NADH	0.033 mM
		(Table 1)
Substrate specificity	: NADH	100 %
	NADPH	1 %
	MTT	100 %
	Lipoate	103 %
		(Table 1)

STORAGE

Store at -20°C


APPLICATION

The enzyme is useful for measurement of various dehydrogenase reactions in the visible spectral range.

ASSAY

Principle

The change in absorbance is measured at 565 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of Diaphorase that forms 1 μmol of NAD^+ per minute at 30 °C

Solutions

- I Buffer solution ; 100 mM HEPES, pH 7.0
- II 3-(4,5-Dimethyl-2-thiazolyl)-2,5-diphenyl-2*H*-tetrazolium bromide (MTT) solution ; 10 mM (20 mg MTT disodium salt·2H₂O/5 mL distilled water)
- III NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O /10 mL distilled water)
- IV Triton solution ; 10 % (1 mL TritonX-100 dilute with distilled water up to 10 mL)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 1 to 5 U/mL with 50 mM potassium phosphate buffer, pH 7.5, 1mg/mL BSA.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	15.00 mL	Solution IV	1.50 mL
Solution II	1.50 mL	H ₂ O	10.80 mL
Solution III	1.20 mL		
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 565 nm per minute (ΔAbs_{565}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{565}) \times (3.00 + 0.01)}{20.0 \times 0.01} \quad \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

20.0 ; millimolar extinction coefficient of MTT ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Packman, L.C., and Perham. R.N.; *FEBS Lett.*, **139**, 155 (1982)

Table 1. SUBSTRATE SPECIFICITY OF DIAPHORASE 22

Acceptor	MTT	Lipoate
Km _{Acceptor} (mM)	0.345	2.0
Km _{NADH} (mM)	0.033	0.01
Relative Activity	100	103
Assay Mixture	HEPES (pH 7.0) 50 mM NADH 0.5 mM MTT 0.5 mM Triton X-100 0.5 %	Potassium Phosphate (pH 6.5) 70.5 mM NADH 0.2 mM NAD 0.3 mM Lipoate 10.2 mM EDTA 0.81 mM BSA 0.7 mg/mL
Wavelength for Measurement (nm)	565	340
Extinction coefficient (cm ² /μmol)	20	6.22




Fig. 1 pH profile

(\triangle Gly-KOH, \circ Bicine,
● phosphate, \blacktriangle HEPES)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following
buffer solution (0.1 M), :
 \triangle Gly-KOH, \circ Bicine,
● phosphate, \blacktriangle HEPES,
 \square MES, \blacksquare Citrate)




Fig. 3 Thermal stability

(treated for 15 min in 0.1M
potassium phosphate buffer,
pH 7.5)




Fig. 4 Thermal stability

(treated for in 0.1M potassium
phosphate buffer, pH 7.5
 \circ 60 °C, \square 70 °C, \bullet 80 °C)

GALACTOSE DEHYDROGENASE (GalDH)

[EC 1. 1. 1. 48]

from recombinant *E. coli*



SPECIFICATION

State	: Ammonium sulphate suspension	
Specific activity	: more than 80 U/mg protein	
Contaminants	: (as GalDH activity = 100 %)	
	NADH oxidase	< 0.10 %
	LDH	< 0.10 %
	ADH	< 0.01 %

PROPERTIES

Subunit molecular weight	: ca. 33,800	
Optimum pH	: 10.5	(Fig. 1)
pH stability	: 5.0 - 10.0	(Fig. 2)
Thermal stability	: No significant decrease in activity up to 50 °C with Ammonium sulphate and 40 °C without Ammonium sulphate .	(Fig. 3, 4)
Michaelis constants	: D-Galactose	0.25 mM
	NAD ⁺	0.15 mM
Substrate specificity (100mM)	: D-Galactose	100 %
	D-Glucose	0.2 %
	D-Xylose	8.7 %
	D-Maltose	0.1 %
	D-Sucrose	0.1 %

STORAGE

Store at 2 to 10 °C (Do not freeze)
Stable at 4 °C for at least one year


APPLICATION

This enzyme is useful for determination of galactose.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of GalDH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH9.1 (at 30 °C)
- II NAD⁺ solution ; 100 mM
- III D-Galactose solution ; 1 M
- IV Enzyme diluent ; 20 mM potassium phosphate, 0.1 % bovine serum albumin, pH 7.5

Preparation of Enzyme Solution

Dilute the enzyme suspension to approx. 5 U/mL with the enzyme diluent.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.
 - Solution I 27.60 mL
 - Solution II 0.90 mL
 - Solution III 1.50 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by the Bradford's method




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacktriangle Glycine-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (0.1
 M); Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacktriangle Glycine-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 25 mM
 potassium phosphate buffer
 pH 7.5, with or without 3.2 M
 ammonium sulphate (AmS).)




Fig. 4 Thermal stability

(treated in 25 mM potassium
 phosphate buffer pH 7.5 at \circ
 40 °C, \square 50 °C, \bullet 60 °C
 without ammonium sulphate.)





Fig. 5 Storage Stability

(ammonium sulphate suspension
(ca. 1300 U/mL) store at 4 °C(○))

GLUCOKINASE 2 (GlcK2)

[EC 2. 7. 1. 2]

from *Recombinant E.coli*

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 350 U/mg protein	
Contaminants	: (as GlcK2 activity = 100 %)	
	Glucose-6-phosphate dehydrogenase	< 0.01 %
	Phosphoglucomutase	< 0.01 %
	6-Phosphogluconate dehydrogenase	< 0.01 %
	Hexose-6-phosphate isomerase	< 0.01 %
	Glutathione reductase	< 0.01 %

PROPERTIES

Subunit molecular weight	: ca. 32,000	
Optimum pH	: 9.0	(Fig. 1)
pH stability	: 7.0 - 10.0	(Fig. 2)
Optimum temperature	: 70 °C	(Fig. 5)
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)
Michaelis constants	: (60 mM Tris-HCl buffer, pH 8.5, at 30 °C)	
	Glucose	0.1 mM
	ATP	0.05 mM
Substrate specificity	: D-Glucose	100 %
	D-Mannose	20 %
	D-Fructose	0 %

STORAGE

Stable at -20 °C for at least one year

APPLICATION

The enzyme is useful for diagnostic reagent, for example, glucose determination or CK determination, and for the specific determination of glucose.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of GlcK2 that forms 1 μmol of glucose-6-phosphate per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH 9.0
- II ATP solution ; 100 mM (0.605 g ATP disodium salt·3H₂O/(8.2 mL distilled water + 1.8 mL 1 N-NaOH))
- III MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O/100 mL distilled water)
- IV NADP⁺ solution ; 22.5 mM [(0.172 g NADP⁺ monosodium salt or 0.177 g NADP⁺ disodium salt)/10 mL distilled water]
- V Glucose solution ; 40 mM (0.072 g glucose (anhyd.)/10 mL distilled water)
- VI Glucose-6-phosphate dehydrogenase (G6PDH) ; (from yeast. Roche Diagnostics K.K., No. 127 671) suspension in 3.2 M (NH₄)₂SO₄ solution (10 mg/2 mL) approx. 140 U/mg at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	17.97 mL	Solution IV	1.20 mL
Solution II	1.20 mL	Solution V	9.00 mL
Solution III	0.60 mL	Solution VI	0.03 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADPH (cm²/μmol)

*Protein concentration; determined by Bradford's method

REFERENCE

1. Hengartner, H., and Zuber, H.; *FEBS Lett.*, **37**, 212 (1973)
2. Kamei, S., Tomita, K., Nagata, K., Okuno, H., Shiraishi, T., Motoyama, A., Ohkubo, A., and Yamanaka, H.; *J. Clin. Biochem. Nutr.*, **3**, 1 (1987)
3. Tomita, K., Kamei, S., Nagata, K., Okuno, H., Shiraishi, T., Motoyama, A., Ohkubo, A., and

Yamanaka, M.; *ibid.*, **3**, 11 (1987)




Fig. 1 pH profile

(
 △ acetate, ○ phosphate,
 ● Tris-HCl, ▲ carbonate
)




Fig. 2 pH stability

(
 treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M), :
 △ acetate, ○ phosphate,
 ● Tris-HCl, ▲ carbonate
)




Fig. 3 Thermal stability

(
 treated for 15 min. in 0.1 M
 Tris-HCl buffer, pH 8.9
)




Fig. 4 Thermal stability

(
 treated in 0.1 M Tris-HCl
 buffer , pH 8.9
 ○ 60 °C, □ 70 °C, ● 80 °C
)





Fig. 5 Thermal activity

(defined as 100 % at 70 °C)

GLUCOSE DEHYDROGENASE (GlcDH2)

[EC 1. 1. 1. 47]

from recombinant *E. coli*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 900 U/mg protein	
Contaminants	: (as GlcDH2 activity = 100 %)	
	NADH oxidase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 126,000	
Subunit molecular weight	: ca. 31,500	
Optimum pH	: 8.5	(Fig. 1)
pH stability	: 5.0 - 10.0 (with 3M NaCl)	(Fig. 2)
Thermal stability	: No significant decrease in activity up to 70 °C. (with 3M NaCl and 0.1% BSA)	(Fig. 3, 4)
Michaelis constants	: D-Glucose	3.7 mM
	NAD ⁺	0.06 mM
	NADP ⁺	0.02 mM
Substrate specificity (100mM)	: D-Glucose	100 %
	D-Maltose	1.1 %
	D-Galactose	0.1 %
	D-Xylose	3.0 %
	D-Fructose	0.3 %
	D-Mannose	4.8 %
	D-Arabinose	0 %
	Trehalose	0 %
	D-Lactose	1.3 %
	D-Sucrose	0 %
	2-Deoxy-D-Glucose	100 %
	D-Glucose-1-Phosphate	0 %
	D-Glucose-6-Phosphate	0 %
	D-Sorbitol	0 %

STORAGE

Stable at -20 °C for at least one year

APPLICATION

This enzyme is useful for determination of glucose.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of GlcDH2 that forms 1 μmol of NADH per minute at 37 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH 8.5 (at 25 °C)
- II NAD⁺ solution ; 100 mM (0.663 g NAD⁺ free acid/10 mL distilled water)
- III D-Glucose solution ; 1 M (1.802 g glucose (anhyd.)/10 mL distilled water)
- IV NaCl solution ; 5 M (2.92 g NaCl/10 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 15 U/mL with 20 mM potassium phosphate buffer containing 1mg/mL BSA and 2 M NaCl, pH 6.5.

Procedure

1. Prepare the following reaction mixture and pipette 2.70 mL of reaction mixture into a cuvette.

Solution I	17.22 mL
Solution II	0.50 mL
Solution III	2.00 mL
Solution IV	0.28 mL
2. Incubate at 37 °C for about 3 minutes.
3. Add 0.015 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (2.70 + 0.015)}{6.22 \times 0.015} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by the absorbance at 280nm (Abs_{280}),
where 1 Abs_{280} = 1 mg/mL

REFERENCE

1. Ramaley, R.F. and Vasantha, N.; *J. Biol. Chem.* **258**, 12558-12565 (1983)




Fig. 1 pH profile

[○ phosphate, ● Tris-HCl, ▲ glycine]




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following
buffer solution (0.1 M) containing 3 M
NaCl : △ acetate, ○ phosphate,
● Tris-HCl, ▲ glycine)




Fig. 3 Thermal stability

(treated for 15 min in 0.1M phosphate
buffer, pH 6.5, containing 3 M NaCl
and 0.1 % BSA)




Fig. 4 Thermal stability

(treated for in 0.1 M phosphate buffer,
pH 6.5, containing 3M NaCl and 0.1 %
BSA, ○ 60°C, □ 70°C, ● 80°C)

D-LACTATE DEHYDROGENASE (D-LDH)

[EC 1. 1. 1. 28]

from *Microorganism*



FOR PYRUVATE → LACTATE REACTION

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 2,500 U/mg protein	
Contaminants	: (as D-LDH activity = 100 %)	
	NADH oxidase	< 0.01 %
	GOT	< 0.01 %
	GPT	< 0.01 %

PROPERTIES


Molecular weight	: ca. 80,000	
Subunit molecular weight	: ca. 40,000	
Optimum pH	: 7.5	(Fig. 1)
pH stability	: 5.5 - 10.0	(Fig. 2)
Isoelectric point	: 4.1	
Thermal stability	: No detectable decrease in activity up to 40 °C.	(Fig. 3, 4)
Michaelis constants	: (94 mM Potassium phosphate buffer, pH 7.5, at 30 °C)	
	Pyruvate	0.80 mM
	NADH	0.18 mM
Stabilizers	: (NH ₄) ₂ SO ₄ , BSA	
Inhibitors	: Zn ²⁺ , Cu ²⁺	

STORAGE

Stable at -20 °C at least one year

ASSAY**Principle**

The change in absorbance is measured at 340 nm according to the following reaction.

**Unit Definition**

One unit is defined as the amount of D-LDH that forms 1 μmol of NAD^+ per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Potassium phosphate buffer, pH 7.5
- II Sodium pyruvate solution ; 100 mM (100 mg sodium pyruvate/10 mL distilled water)
- III NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 3 to 5 U/mL with 50 mM potassium phosphate buffer containing 1 mg/mL BSA, pH 7.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.
Solution I 28.00 mL
Solution II 1.20 mL
Solution III 0.80 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method




Fig. 1 pH profile

(Δ acetate, \circ phosphate,)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
 Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M potassium phosphate buffer, pH 7.0)




Fig. 4 Thermal stability

(treated in 0.1 M potassium phosphate buffer, pH 7.0
 \circ 40 °C, \square 45 °C, \bullet 50 °C)

MALATE DEHYDROGENASE (MDH)

[EC 1. 1. 1. 37]

from *Microorganism*



FOR OXALATE → MALATE REACTION

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 1,200 U/mg protein	
Contaminants	: (as MDH activity = 100 %)	
	GOT	< 0.01 %
	GPT	< 0.01 %
	NADHoxidase	< 0.01 %
	Glutamate dehydrogenase	< 0.01 %
	Fumarase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 72,000	
Subunit molecular weight	: ca. 36,000	
Optimum pH	: 9.0	(Fig. 1)
pH stability	: 5.5 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 50 °C.	(Fig. 3, 4)
Michaelis constants	: (90 mM Tris-HCl buffer, pH 9.0, at 30 °C)	
	Oxaloacetate	0.027 mM
	NADH	0.014 mM

STORAGE

Stable at -20 °C for at least six months


APPLICATION

This enzyme is useful for enzymatic determination of L- malate and of glutamate oxaloacetate transaminase in clinical analysis.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of MDH that forms 1 μmol of NAD^+ per minute at 30 °C.

Solutions

- I Buffer solution ; 200 mM Tris-HCl, pH 9.0
- II Oxaloacetate solution ; 15 mM (0.020 g oxaloacetate/10 mL distilled water)
- III NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 3 to 5 U/mL with 100 mM Tris-HCl buffer, pH 9.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	13.50 mL
Solution II	1.00 mL
Solution III	0.57 mL
H ₂ O	14.93 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method




Fig. 1 pH profile

(Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M);
 Δ acetate, \circ phosphate,
 \bullet Tris-HCl, \blacksquare Gly-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M
 Tris-HCl buffer, pH 9.0)




Fig. 4 Thermal stability

(treated in 0.1 M Tris-HCl
 buffer, pH 9.0
 \circ 50 °C, \square 55 °C, \bullet 60 °C)

MUTAROTASE (MRO)

[EC 5. 1. 3. 3]

from *Microorganism*

α -D-Glucose \leftrightarrow β -D-Glucose

SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 120 U/mg protein	
Contaminants	: (as MRO activity = 100 %)	
	NADHoxidase	< 0.01 %

PROPERTIES

Subunit molecular weight	: ca. 39,500	
Optimum pH	: 7.0 - 9.0	(Fig. 1)
pH stability	: 3.5 - 10.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 50 °C.	(Fig. 3, 4)

STORAGE

Stable at -20 °C for at least one year


APPLICATION

This enzyme is useful for enzymatic determination of glucose.

ASSAY

Principle

Acceleration of the glucose dehydrogenase reaction by Mutarotase is measured according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of Mutarotase that forms 10 μmol of NADH per minute at 25 $^{\circ}\text{C}$.

Solutions

- I HEPES buffer ; 50 mM (1.19 g HEPES / 100 mL distilled water, adjust pH to 7.5 with NaOH)
- II Sucrose solution ; 16.7 mM (57 mg Sucrose / 10 mL distilled water)
- III NAD^+ solution ; 100 mM (0.663 g NAD^+ free acid / 10 mL distilled water)
- IV Glucose dehydrogenase solution ; 3 kU/mL (GlcDH2, Nipro Corp. / 20 mM potassium phosphate containing 2 M NaCl, pH6.5)
- V β Fructosidase solution ; ≥ 30 kU/mL (100 mg Invertase from baker's yeast, Sigma-Aldrich I4504 / 1 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 0.7 to 1.4 U/mL with 20 mM potassium phosphate pH 7.3 containing 1 mg/mL BSA.

Procedure

1. Prepare the following reaction mixture and pipette 2.70 mL of reaction mixture into a cuvette.

Solution I	19.90 mL	Solution IV	0.166 mL
Solution II	1.00 mL		
Solution III	0.60 mL		
2. Add 0.015 mL of the enzyme solution into the cuvette and mix.
3. Incubate at 25 $^{\circ}\text{C}$ for about 3 minutes.
4. Add 0.06 mL of the Solution V into the cuvette and mix.
5. Read absorbance change at 340 nm per minute (ΔAbs1) in the linear portion of curve.
6. Run the procedure 1 to 5 with the enzyme diluent instead of the enzyme solution (ΔAbs2).

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs1} - \Delta\text{Abs2}) \times (2.70 + 0.015 + 0.060)}{6.22 \times 0.015 \times 10} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

10 ; conversion factor

*Protein concentration ; determined by the absorbance at 280 nm (Abs_{280}),
where 1 $\text{Abs}_{280} = 1 \text{ mg/mL}$




Fig. 1 pH profile

(
 △ acetate, □ MES, ◆ PIPES,
 ◇ HEPES, ● Tris-HCl,
 ■ Bicine, ▲ Glycine-KOH
)




Fig. 2 pH stability

(
 treated for 24 hr at 4 °C in the
 following buffer solution (0.1 M)
 containing 0.1 % BSA;
 △ acetate, ○ phosphate,
 ● Tris-HCl, ▲ Glycine-KOH
)




Fig. 3 Thermal stability

(
 treated for 15 min in 0.1 M potassium
 phosphate buffer, pH 6.5, 0.1 % BSA.
)




Fig. 4 Thermal stability

(
 treated in 0.1 M potassium
 phosphate buffer, pH 6.5, 0.1 % BSA
 at ○ 40 °C, □ 50 °C, ● 60 °C
)

PHENYLALANINE DEHYDROGENASE (PheDH)

[EC 1.4.1.20]

from *Thermoactinomyces intermedius*

SPECIFICATION

State	: Ammonium sulphate suspension	
Specific activity	: more than 30 U/mg protein	
Contaminants	: (as PheDH activity = 100 %)	
	NADH oxidase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 380,000	
Subunit molecular weight	: ca. 40,000	
Optimum pH	: 11.5	(Fig. 1)
pH stability	: 5.0 - 10.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 50 °C.	(Fig. 3, 4)
Michaelis constants	: (200 mM Gly-KCl-KOH buffer, pH 11.0, at 30 °C)	
	L-Phenylalanine	0.66 mM
	NAD ⁺	0.05 mM
Substrate specificity	: L-Phenylalanine	100 %
	L-Tyrosine	7.6 %
	L-Methionine	1.5 %


STORAGE

Stable at 4 °C for at least six months (Do not freeze)

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of PheDH that forms 1 μmol of NADH per minute at 30 °C.

Solutions

- I Buffer solution ; 400 mM Gly-KCl-KOH, pH 11.0
- II L-Phenylalanine solution ; 100 mM (0.165 g L-phenylalanine/10 mL distilled water)
- III NAD^+ solution ; 100 mM (0.663 g NAD^+ free acid/10 mL distilled water)

Preparation of Enzyme Solution

Dilute the ammonium sulphate suspension of enzyme to 2 to 6 U/mL with 10 mM Tris-HCl buffer, pH 8.0, containing 50 mM NaCl.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	15.00 mL
Solution II	3.00 mL
Solution III	0.15 mL
H_2O	11.85 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 340nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method

REFERENCE

1. Ohshima, T., Takada, H., Yoshimura, T., Esaki, N., and Soda, K.; *J. Bacteriol.*, **173**, 3943 (1991)




Fig. 1 pH profile

■ Gly-KCl-KOH




Fig. 2 pH stability

treated for 24 hr at 4 °C in the following buffer solution (50 mM);
 △ acetate, ○ phosphate,
 ■ Gly-KCl-KOH




Fig. 3 Thermal stability

treated for 15 min in 10 mM potassium phosphate buffer, pH 7.2




Fig. 4 Thermal stability

treated in 10 mM potassium phosphate buffer, pH 7.2
 ○ 50 °C, □ 60 °C, ● 70 °C

PHOSPHOTRANSACETYLASE (PTA2)

[EC 2. 3. 1. 8]

from recombinant *E. coli*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 5,000 U/mg protein	
Contaminants	: (as PTA2 activity = 100 %)	
	Acetate kinase	< 0.01 %
	Adenylate kinase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Molecular weight	: ca. 69,700	
Subunit molecular weight	: ca. 33,500	
Optimum pH	: 7.4	(Fig. 1)
pH stability	: 4.0 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)
Michaelis constants	: (87mM Tris-HCl buffer, pH 7.5, at 30 °C)	
	Coenzyme A	0.1 mM
	Acetyl Phosphate	0.5 mM

STORAGE

Stable at -20 °C for at least one year

ASSAY

Principle

The change in absorbance is measured at 233 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of PTA that forms 1 μmol of acetyl-CoA per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Tris-HCl, pH 7.5
- II CoA solution ; 6.4 mM (50 mg CoA trilithium salt/10 mL distilled water)
- III Acetylphosphate solution ; 217 mM (0.400 g acetylphosphate potassium lithium salt/10 mL distilled water)
- IV Ammonium sulfate (AmS) solution ; 1 M (13.2 g AmS/100 mL distilled water)

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 20 U/mL with 50 mM Tris-HCl buffer, pH 8.0.

Procedure

1. Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	26.0 mL	Solution III	1.0 mL
Solution II	2.0 mL	Solution IV	1.0 mL
2. Incubate at 30 °C for about 3 minutes.
3. Add 0.01 mL of enzyme solution into the cuvette and mix.
4. Read absorbance change at 233 nm per minute (ΔAbs_{233}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{233}) \times (3.00 + 0.01)}{4.44 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

4.44 ; differential millimolar extinction coefficient between acetyl-CoA and CoA ($\text{cm}^2/\mu\text{mol}$)

*Protein concentration ; determined by Bradford's method




Fig. 1 pH profile

(
 △ acetate, ○ phosphate,
 ● Tris-HCl, ■ Gly-KOH
)




Fig. 2 pH stability

(
 treated for 24 hr at 4 °C in the
 following buffer solution (50 mM);
 △ acetate, ○ phosphate,
 ● Tris-HCl, ■ Glycine-KOH
)




Fig. 3 Thermal stability

(
 treated for 15 min in 50 mM
 Tris-HCl buffer, pH 8.5,
)





Fig. 4 Thermal stability

(
 treated in 50 mM Tris-HCl
 buffer, pH 8.5, 0.1 % BSA
 ○ 55 °C, □ 60 °C, ● 65 °C
)

POLYNUCLEOTIDE PHOSPHORYLASE (PNPase3)

[EC 2. 7. 7. 8]

from recombinant *E. coli*



FOR DEPOLYMERIZATION REACTION

SPECIFICATION

State	: Liquid
Specific activity	: more than 2,000 U/mg protein

PROPERTIES

Subunit molecular weight	: ca. 79,000	
Optimum pH	: 9.0 - 9.5	(Fig. 1)
pH stability	: 8.0 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 55 °C.	(Fig. 3, 4)
Effectors	: cations and anions	(Fig. 5, 6)

STORAGE

at -20 °C



APPLICATION

The enzyme is useful for the preparation of polyribonucleotide.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reactions.



Unit Definition

One unit of activity is defined as the amount of PNPase that forms 1 μmol of ADP per hour at 60 °C by depolymerizing of Poly A.

Solutions

(Reaction I)

- I Buffer solution ; 100 mM Tris-HCl, pH 9.5 ((1.212 g Tris + 0.074 g EDTA + 0.014 mL 2-mercaptoethanol + 0.610 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ + 0.746 g KCl)/80 mL distilled water, adjusted to pH 9.5 with 1 N-HCl and filled up to 100 mL with distilled water)
- II KH_2PO_4 solution ; 65 mM (0.088 g KH_2PO_4 /10 mL distilled water)
- III polyadenylate (Poly A) solution ; (25 mg Poly A potassium salt/1 mL distilled water; ca. 35 mM based on AMP concentration)

(Reaction II)

- IV Buffer solution ; 100 mM Triethanolamine buffer, pH 7.6 ((9.300 g triethanolamine-HCl + 0.407 g $\text{MgCl}_2 \cdot 6\text{H}_2\text{O}$ + 0.373 g KCl)/400 mL distilled water, adjusted to pH 7.6 with 1 N-NaOH and filled up to 500 mL with distilled water)
- V NADH solution ; 13.1 mM (0.100 g NADH disodium salt $\cdot 3\text{H}_2\text{O}$ /10 mL distilled water)
- VI Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- VII Pyruvate kinase (PK) ; (from rabbit muscle, Roche Diagnostics K.K., No. 128 155) crystalline suspension in 3.2 M $(\text{NH}_4)_2\text{SO}_4$ solution (10 mg/mL) approx. 200 U/mg at 25 °C
- VIII Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., Product Code: LDH-02) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 2 to 150 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

(Reaction I)

1. Prepare the following reaction mixture and pipette 0.55 mL of reaction mixture into a test tube.

Solution I	2.50 mL	Solution III	1.00 mL
Solution II	1.00 mL	H ₂ O	1.00 mL
 2. Add 0.10 mL of enzyme solution and mix.
 3. Incubate at 60 °C for exactly 10 minutes.
 4. After incubation, add 0.01 mL conc. HCl and mix.
 5. Centrifuge at 10,000 rpm for 30 seconds.
- At the same time, repeat the Procedure 1 to 5 using distilled water in place of enzyme solution in Procedure 2 (as blank).

(Reaction II)

6. Prepare the following reaction mixture and pipette 2.50 mL of the reaction mixture into a cuvette.

Solution IV	24.18 mL	Solution VII	0.12 mL
-------------	----------	--------------	---------

Solution V 0.40 mL

Solution VIII 0.05 mL

Solution VI 0.25 mL

7. Incubate at 30 °C for about 3 minutes.
 8. Add 0.10 mL of supernatant of Procedure 5 and mix.
 9. Read absorbance at 340 nm (Abs•test).
- Repeat the Procedure using blank (Abs•blank).

Calculation

$$\text{Volume activity (U/mL)} = ((\text{Abs} \cdot \text{blank}) - (\text{Abs} \cdot \text{test})) \times \frac{2.60 \times 0.65}{6.22 \times 0.10 \times 0.10} \times \frac{60}{10} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH (cm²/μmol)

*Protein concentration ; determined by the absorbance at 280nm (Abs₂₈₀),
where 1 Abs₂₈₀ = 1 mg/mL

REFERENCES

1. Smith, J.C., and Eaton, M.A.W.; *Nucleic Acids Research*, **1**, 1763 (1974)
2. Wood, J.N., and Hutchinson, D.W.; *ibid.*, **3**, 219 (1976)




Fig. 1 pH profile

(● Tris-HCl)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the following buffer solution (0.1 M);
● Tris-HCl, ■ Glycine-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 0.1 M Tris-HCl buffer, pH 8.5)




Fig. 4 Thermal stability

(treated in 50 mM Tris-HCl buffer, pH 8.5
○ 55 °C, □ 60 °C, ● 65 °C)




Fig. 5 Effect of various cations on the activity of PNPase3 in the following Assay

Measurement: 0.015 mL of each cation solution, 0.010 mL of enzyme solution and 0.055 mL of reaction mixture were mixed, and reacted at 60 °C. After 10 minutes, the quantity of ADP was determined.
○ NaCl, △ KCl, □ MgCl₂





Fig. 6 Effect of various anions on the activity of PNPase3 in the following Assay

Measurement: 0.015 mL of each anion solution, 0.010 mL of enzyme solution and 0.055 mL of reaction mixture were mixed, and reacted at 60 °C. After 10 minutes, the quantity of ADP was determined.
○ NaCl, △ CH₃COONa, □ Na₂SO₄, ● NaH₂PO₄

PYRUVATE KINASE (PK2)

[EC 2.7.1.40]

from Recombinant *E. coli*



SPECIFICATION

State	: Lyophilized	
Specific activity	: more than 230 U/mg protein	
Contaminants	: (as PK activity = 100 %)	
	Adenylate kinase	< 0.01 %
	Lactate dehydrogenase	< 0.01 %

PROPERTIES

Subunit molecular weight	: ca. 60,000	
Optimum pH	: 6.0 - 6.5	(Fig. 1)
pH stability	: 6.0 - 11.0	(Fig. 2)
Thermal stability	: No detectable decrease in activity up to 60 °C.	(Fig. 3, 4)
Michaelis constants	: (76 mM Imidazole-HCl buffer, pH 7.2, at 30 °C)	
	Phosphoenolpyruvate	1.0 mM
	ADP	1.5 mM

STORAGE

Stable at -20 °C for at least one year


APPLICATION

The enzyme is useful for diagnostic reagent, for example, ADP determination.

ASSAY

Principle

The change in absorbance is measured at 340 nm according to the following reaction.



Unit Definition

One unit of activity is defined as the amount of PK2 that forms 1 μmol of pyruvate per minute at 30 °C.

Solutions

- I Buffer solution ; 100 mM Imidazole-HCl, pH 7.2
- II ADP solution ; 100 mM (0.45 g ADP sodium salt, Sigma-Aldrich A2754/(9.0 mL distilled water + 1.0 mL 1 N NaOH))
- III NADH solution ; 13.1 mM (0.100 g NADH disodium salt·3H₂O/10 mL distilled water)
- IV Phosphoenolpyruvate (PEP) solution ; 56 mM (0.150 g PEP MCA salt/10 mL distilled water)
- V MgCl₂ solution ; 1 M (20.33 g MgCl₂·6H₂O/100 mL distilled water)
- VI KCl solution ; 2.5 M (18.64 g KCl/100 mL distilled water)
- VII Lactate dehydrogenase (LDH) ; (from pig heart, Oriental Yeast Co. Ltd., LDH (P.H.)) ammonium sulfate suspension, approx. 5,000 U/mL at 25 °C

Preparation of Enzyme Solution

Dissolve the lyophilized enzyme with distilled water and dilute to 5 to 10 U/mL with 50 mM Tris-HCl buffer, pH 8.5.

Procedure

- Prepare the following reaction mixture and pipette 3.00 mL of reaction mixture into a cuvette.

Solution I	22.71 mL	Solution V	0.48 mL
Solution II	2.40 mL	Solution VI	0.90 mL
Solution III	0.45 mL	Solution VII	0.06 mL
Solution IV	3.00 mL		
- Incubate at 30 °C for about 3 minutes.
- Add 0.01 mL of enzyme solution into the cuvette and mix.
- Read absorbance change at 340 nm per minute (ΔAbs_{340}) in the linear portion of curve.

Calculation

$$\text{Volume activity (U/mL)} = \frac{(\Delta\text{Abs}_{340}) \times (3.00 + 0.01)}{6.22 \times 0.01} \times \text{d.f.}$$

$$\text{Specific activity (U/mg protein)} = \frac{\text{Volume activity (U/mL)}}{\text{Protein concentration (mg/mL)}^*}$$

d.f. ; dilution factor

6.22 ; millimolar extinction coefficient of NADH (cm²/μmol)

*Protein concentration ; determined by Bradford's method

REFERENCE

- Sakai, H., Suzuki, K., and Imahori, K.; *J. Biochem.*, **99**, 1157 (1986)




Fig. 1 pH profile

(Δ Acetate, \square Imidazole-HCl,
 \bullet Tris-HCl, \blacksquare Glycine-KOH)




Fig. 2 pH stability

(treated for 24 hr at 4 °C in the
 following buffer solution (50 mM);
 Δ Acetate, \square Imidazole-HCl,
 \bullet Tris-HCl, \blacksquare Glycine-KOH)




Fig. 3 Thermal stability

(treated for 15 min in 50 mM
 Tris-HCl buffer, pH 8.5)




Fig. 4 Thermal stability

(treated in 50 mM Tris-HCl
 buffer, pH 8.5
 \circ 55 °C, \square 60 °C, \bullet 65 °C)